

REZENOPY™ (reh zeh' noe pee)
(naloxone HCl) nasal spray
10 mg

QUICK START GUIDE

Use REZENOPY nasal spray for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

Do not unpack REZENOPY nasal spray until ready to use.

Do not test REZENOPY nasal spray before use.

Each REZENOPY nasal spray device contains 1 dose of medicine and cannot be reused.

1

Check person for signs of suspected opioid overdose.


Shout “wake up” and **shake** the person gently.

- unresponsive
- unconscious or unable to wake up
- slow, shallow, irregular breathing
- fingernails or lips turning blue or purple




2

Lay the person flat on their back.



3


Peel open package and remove REZENOPY nasal spray from box.



4

Hold REZENOPY nasal spray as shown with your index and middle fingers on either side of the nozzle and your thumb on the bottom of the plunger.

Do not apply pressure until you are ready to give the dose.



5

Provide support behind the neck with your hand and tilt the person's head back.




Gently insert the nozzle tip all the way into one nostril.



6

Push the plunger hard and all the way in to give the dose.

Remove REZENOPY nasal spray from nostril after the dose has been given.



7

Call for 911 or get emergency help right away.




8

Put the person into the recovery position by turning them on their side.

Watch the person closely for **2 to 3 Minutes**




Put in Recovery Position and Watch



9

If the person does not respond, get a new REZENOPY nasal spray from the box, and give another dose into the **other nostril**.



If more REZENOPY nasal spray is available, repeat steps 2 through 6 every 2 to 3 minutes, alternating nostrils, until the person wakes up or medical help arrives. Stay with the person until medical help arrives.



10

Throw Away the used REZENOPY nasal spray in a place away from children.

